

Global Agronomy Research Journal

Decontaminating Test of Dadih Against Pathogenic Microorganisms and Allure Belongings on Native Poultry Privilege Means

Nurzainah Ginting

Animal Science Study Program, Faculty of Agriculture, Universitas Sumatera Utara, Medan 20155, Indonesia

* Corresponding Author: Nurzainah Ginting

Article Info

ISSN (online): 3049-0588

Volume: 01

Issue: 04

July-August 2024

Received: 15-06-2024

Accepted: 05-07-2024

Page No: 01-03

Abstract

Dadih holds lactic acid microorganisms that are antimicrobial because it maybe appropriated to assert the well-being of fowl to a degree native chickens. This study aims to demonstrate the skill of dadih to restrain the progress of pathogenic microorganisms, that is to say Staphylococcus aureus, Escherichia coli and Poisoning caused by eating food. This study will likewise search for out invivo effect of dadih on the means privilege of native poultry contaminated accompanying Poisoning caused by eating food. Antimicrobial research uses plate spread system accompanying antimicrobial district limit. The invivo study was transported utilizing a entirely randomized design accompanying 4 situations and five replications place the limit was the growth of privilege means that is to say organ meat, bag fabrisius and hate. The results revealed that the antimicrobial test of dadih against Staphylococcus aureus, Escherichia coli and Poisoning caused by eating food made a district of restriction of 7.84, 6.63 and 8.34 mm. Dadih situation considerably distressed the portion of organ meat, bag fabrisius and hate. As end that dadih is smart to restrain the tumor of Staphylococcus aureus, Escherichia coli and Poisoning caused by eating food at medium substance accompanying a range of restriction district middle from two points 6.63-8.34. Dadih is likewise intelligent to claim the well-being of native hen.

Keywords: medicine, bag fabrisius, dadih, native poultry, hate, organ meat

1. Introduction

The Economic unit dictated a ban on the use of medicines as feed supplements. Medicines can likewise not any more be used to fix development or weak cleanliness. This is on account of likely concerns concerning their attainable incorporation in animals raised on a farm amount. Moreover, opposition in mammals maybe communicated to persons and harm to their fitness. Irregularities to the use of medicines are admitted to treat cases of afflictions ^[10, 2]. The occupancy of medicine residues in gist can have disadvantageous belongings. In Denmark, earlier, they secondhand over 105 rhythmical tonnes of medicines for tumor publicity but now the medicine use is ^[3].

Individual of the potential options for the happening of alternative medicines search out use probiotics that are organic fabrics to boost the invulnerable order of animals raised on a farm. Individual model of probiotics is confuse milk dadih. Nation in West Sumatra eat dadih that they calls dadih as a side plate and established cure. Dadih is created from a usual effervescence process that uses bamboo tubes as cartons and joker leaves as covers. The effervescence process can happen on account of the vicinity of microorganisms on the bamboo division and likewise on the central surface of the bamboo hose ^[4]. Dadih holds plenty lactic acid microorganisms that is about 75%, e.g., Lactobacillus and Bifidobacterium ^[5].

An sour pH of dadih has the talent to restrain microbial progress. Also, dadih holds lactic acid microorganisms that produce bacteriocin that can restrict pathogenic bacterial strains in the way that Listeria, Clostridium, and Poisoning caused by eating food. Bacteriocins are cationic peptides that cause container cessation of pathogenic microorganisms by making pores and discharging the cytosolic constituents of pathogenic bacterial containers. As long as, bacteriocins are smart to adjust host privilege with reconstructing host strength ^[6]. Bacteriocins are antimicrobial peptides caused by lactic acid microorganisms that mainly suppres or cancel pathogenic microorganisms raw spot and change the arrangement of the gut microbiota in animal models in the way that rodent, boars, and chickens ^[7, 8].

This study secondhand invitro and *in vivo* tests. The artificial test was administered to test the uncontaminated capacity of dadih against pathogenic microorganisms while the invivo test search out test the effect of dadih devouring on the strength tools of native chickens that is to say organ meat, bag fabrisius and hate. Invitro test utilizing plate spread pattern and calculation of hindrance/completely clean district. Clean paper discs were trickled accompanying 20 micron liters of dadih and admitted to signify nearly 3 hours. Bacterial smearing was acted utilizing a unproductive cotton-tipped stick superficial of MHA radio accompanying bacterial delay in 0.5 McFarland turbidity. Dadih-discussed paper discs were established into the agar that had happened striate accompanying bacterial delay. Hatched for 18-24 hours. The decontaminating endeavor of the dadih was noticed and the width of the clear district made about the paper plate was calculated as decontaminating district.

In the *in vivo* study, following in position or time the chickens were killed, the means that indicate the fitness of the poultry physique were painstakingly captured and weighed, containing the hate, organ meat, and bag fabrisius utilizing examining scales. In addition, each sample was detached for one live burden of the hen manifolded by 100. Invivo test was transported on poultry. Research design secondhand Non-factorial Entirely Randomized Design (CRD) amounting to 4 (four) situations and 5 (five) replications, place: P0 = Control outside dadih presidency; P1 = 25% dadih additional accompanying 75% aquadest; P2 = 50% dadih additional accompanying 50% aquadest; P3 = COLIMAS (medicine). The situation was completed activity through presidency in gulping water. The research dossier got will be analysed utilizing study of difference (ANOVA) and if the results acquired are very legitimate or evident before persisted by utilizing Duncan's Diversified Range Test.

3. Results and Dispute

3.1. Completely clean Test

Antimicrobial district is a limit of dadih's talent to restrain or destroy pathogenic microorganisms. Four catagories of the talent to cancel microorganisms are very forceful if the completely clean district >20 mm, forceful 10-20 mm, medium 5-10 mm and feeble <5 mm^[9, 10] erect that width of district of restriction of Lactobacillus categorized from 9 ± 0.0 to 27 ± 2.83 mm. Lactobacillus has the capacity to restrict Poisoning caused by eating food. In this place study, it was showed that dadih was intelligent to restrain the progress of Staphylococcus aureus, Ecscherechia coli and Poisoning caused by eating food at medium type while dadih has the capital decontaminating district on Poisoning caused by eating food. This skill of dadih will benefit the host, in this place case the native hen. Native chickens observed in country regions and admitted to meander in the surroundings will be more surely polluted accompanying pathogenic microorganisms. Thus, the routine augmenting of dadih causes the chickens to have substance against specific bacterial contaminations. Salmonellosis is individual of the big bacterial contaminations moving monetary fowl, inducing misfortunes in fowl result. Chicken typhoid and pullorum ailment, that are extensive septicemic afflictions, are made by *S. gallinarum* and *S. pullorum* individually and pollute chiefly chickens and turkeys^[11]. Salmonellosis is very damaging to the bovine animals manufacturing. But an even bad importance is that persons the one expend hen gist adulterated accompanying Poisoning caused by eating food

knowledge weighty well-being questions^[12].

Table 1: Completely clean Check Come from of Dadih ahead of Staphylococcus aureus, Escherichia coli and Poisoning caused by eating food

Parameter	Antibacterial zone (mm)
Staphylococcus aureus	7.84
Escherichia coli	6.63
Salmonella	8.34

Akanbi and others. established that *E. coli* contamination causes fowl to exhibit generative lesions that compares accompanying diminished cell result. *E. coli* more causes colibacillosis in fowl flocks^[13].

In the air colibacillosis causes septicaemia (colisepticaemia), pneumonia, egg yolk-cyst contamination, omphalitis, cellulitis^[14]. *S. aureus* is individual of ultimate universal pathogens in the direction of fowl farms. This germ causes asdermatitis, arthritis, osteomyelitis, synovitis, teno-synovitis, femoral head loss, botch-nadir, and comphalitis^[15]. Still, what is not likely is adulteration from fowl core polluted accompanying *S. aureus* namely nibbled by persons creating the poison it produces, that is to say Enterotoxigenic creating foodborne contaminate and poisonous shock disease^[16, 17].

3.2. Invulnerable Tool Allotment of Native Hen

3.2.1. Organ meat

Organ meat portion is displayed in "Table 2". Feed supports foods for the progress and incident of basic lymphoid means, that is to say organ meat, bag fabricius and subordinate lymphoid tools, that is to say the hate^[18]. Feed holding probiotics will provoke the poultry's invulnerable whole because it can control ailment. The clue of the effect concerning this wonder is the diameter/allotment of lymphoid means to a degree organ meat. Research administered by^[19] establish the allotment of organ meat betwixt 0.24 to 0.28. This study determines probiotics Bacterium sp. Research attended by^[20] accompanying the situation of bestowing lactic acid microorganisms-located synbiotic rations establish the portion of organ meat about 0.102 to 0.128. Research by^[21] raise the portion of organ meat in the range of 0.22 to 0.24 accompanying the situation of augmenting contain Lactobacillus sp In this place study, it was visualized that the allotment of bag was better accompanying the supplying of dadih. High-quality bag allotment happen accompanying 50% dadih and this situation was equivalent to the definite control, that is to say the situation of Colimas medicines. Research transported by^[20] accompanying the situation of lactic acid microorganisms-located synbiotic rations erect the allotment of bag about 0.2 to 0.3. Research by^[21] establish the portion of bag in the range of 0.11 to 0.14 accompanying the situation of Lactobacillus sp. This study is better than research result by^[21] cause in this place study dadih was likely, that is a accumulation of differing lactic acid microorganisms and fungi.^[23] established that probiotics can influence whole of the invulnerable whole through antibodies hidden by B lymphocytes and T lymphocytes, happening in an increase in the relative pressure of invulnerable means. In this place study, skilled was a meaningful dissimilarity middle from two points the situation outside dadih and the situation accompanying dadih.

3.2.3. Hate

The content of the hate tool is affected for one supplying of

probiotics and feed vitamins. Bestowing probiotics to the feed skilled is an increase in agent for negating the effect of an infection or poison result guide the length of the hate tool. Feed or vitamins considerably influence the incident of lymphoid tools ^[24]. Research transported by ^[19] erect % hate 0.118 to 0.124 place the situation concerning this study was the supplying of probiotics *Bacterium* sp. In this place study, the portion of hate was better than ^[19]. This can develop a better difference of lactic acid microorganisms precipitating better hate incident. The allotment of hate in this place study is inside the range of hate length that in accordance with ^[24] the rational length of grill hate ranges from 0.10-0.23% of live burden. The breadth of hate bestowing 50% dadih is nearly the maximum content of 0.18%.

4. Conclusion and Recommendation

Dadih is capable to uphold the energy of fowl and this is maybe discovered accompanying the happening of hen strength tool signs that is to say organ meat, bag fabrisius and hate in spite of the poultry was contaminated by *Salmonella*. In addition, dadih maybe substantiated expected smart to restrain the incident of *Salmonella* as proved for one antimicrobial test that got an antimicrobial district of 8.34. Dadih is still capable to restrain the growth of *S. aureus* and *E. coli* as proved for one establishment of an antimicrobial district. The exercise of dadih will benefit fowl farms while caring for persons from contamination accompanying pathogenic microorganisms of fowl inception. Given that making dadih is smooth than insulating lactic acid microorganisms, it hopeful more adept to use dadih in fowl farm.

5. Acknowledgments

This research was financed financially by 2023 Talenta Program Universitas Sumatera Utara, Indonesia.

6. References

1. Jerab J, Jansen W, Blackwell J, Van Hout J, Palzer A, Lister S, *et al.* Real-experience dossier on antimicrobial use in European bovine animals: operators, environments, and opportunities. *Antibiotics*. 2022;11(8):1046.
2. Busch G, Kassas B, Palma MA, Risius A. Perceptions of antimicrobial use in livestock breeding and care in Germany, Italy, and the United States. *Livest Sci*. 2020;241:104251.
3. Casewell M, Eriis C, Marco E, McMullin P, Phillips I. The European ban on growth-promoting antibiotics and emerging consequences for human and animal health. *J Antimicrob Chemother*. 2003;52(2):159-161.
4. Yanti SD, Mirwandhono E, Ginting N. Nutrition content of Murrah buffalo's milk curd using lime juice (*Citrus aurantifolia*). *J Peternak Integratif*. 2022;10(2):78-84.
5. Sirait CH, Cahyadi N, Panggabean T, Putu IG. Identifikasi dan pembiakan kultur bakteri pengolah dadih. Laporan Penelitian Balai Penelitian Ternak, Ciawi, Bogor; c1995.
6. Tiwari SK. Bacteriocin-bearing probiotic lactic acid bacteria in controlling dysbiosis of gut microbiota. *Front Cell Infect Microbiol*. 2022;12:851140.
7. Gillor O, Etzion A, Riley MA. The dual role of bacteriocins as anti-competitors and probiotics. *Appl Microbiol Biotechnol*. 2008;81:591-606.
8. Yang SC, Lin CH, Sung CT, Fang JY. Antibacterial

activities of bacteriocins: application in food and pharmaceuticals. *Front Microbiol*. 2014;5:91530.

9. Davis WW, Stout TR. Disc plate method of microbiological antibiotic assay: I. Factors influencing variability and error. *Appl Microbiol*. 1971;22(4):659-65.
10. Dec M, Puchalski A, Nowaczek A, Wernicki A. Antimicrobial activity of *Lactobacillus* strains of poultry origin against bacterial pathogens. *Int Microbiol*. 2016;19(1):57-67.
11. Nabil NM, Tawakol MM, Hassan HM. Assessing the impact of bacteriophages in controlling *Salmonella* in grilled chicken. *Infect Ecol Epidemiol*. 2023;8(1):1539056.
12. Shaji S, Selvaraj RK, Shanmugasundaram R. Enhancement of rice growth and yield in acid sulfate soils by effective acid-tolerant *Rhodopseudomonas palustris* strains for sustainable rice production. *Plant Soil*. 2022;11(11):2814.
13. Logue CM, Wannemuehler Y, Doetkott C, Barbieri NL. Comparative analysis of phylogenetic assignment of human and avian ExPEC and fecal commensal *Escherichia coli* using the revised Clermont phylogrouping method and its correlation with avian pathogenic *Escherichia coli* (APEC) classification. *Front Microbiol*. 2017;8:248958.